

Dipolar dark matter in light of 3.5 keV X-ray Line, Neutrino mass and LUX data

Sudhanwa Patra¹, Nirakar Sahoo² and Narendra Sahu^{2*}

¹ *Centre of Excellence in Theoretical and Mathematical Sciences,
Siksha 'O'Anusandhan University, Bhubaneswar, India*

² *Department of Physics, Indian Institute of Technology,
Hyderabad, Yeddumailaram, 502205, Telengana India*

Abstract

A simple extension of the standard model (SM) providing transient magnetic moments to right-handed neutrinos is presented. In this model, the decay of next-to-lightest right-handed heavy neutrino to the lightest one and a photon ($N_2 \rightarrow N_1 + \gamma$) can explain the ~ 3.5 keV X-ray line signal observed by XMM-Newton X-ray observatory. Beside the SM particles and heavy right-handed Majorana neutrinos, the model contains a singly charged scalar (H^\pm) and an extra Higgs doublet (Σ). Within this minimal set of extra fields the sub-eV masses of left-handed neutrinos are also explained. Moreover, we show that the spin-independent DM-nucleon cross-section is compatible with latest LUX data.

PACS numbers:

*Electronic address: sudha.astro@gmail.com, ph13p1005@iith.ac.in, nsahu@iith.ac.in

I. INTRODUCTION

The existence of dark matter, which constitutes around 26.8% of the total energy budget of the Universe [1], is currently supported by the galaxy rotation curve, gravitational lensing and large scale structure of the Universe [2]. These evidences only indicate the gravitational interaction of dark matter. The detection of dark matter is yet a mystery. The only property of dark matter known to us is its relic abundance and is given by $\Omega_{\text{DM}}h^2 = 0.119$ [1]. Within the framework of the standard model (SM) it is impossible to accomodate a particle candidate of dark matter.

Another aspect of SM is that it does not explain the non-zero neutrino mass which is confirmed by the phenomenon of neutrino oscillation observed in solar, atmospheric and reactor experiments [3]. This indicates towards new physics operative beyond electroweak scale since the SM of particle physics can not explain the same without introducing any extra ingredients. It is found that seesaw mechanisms are the most elegant scenarios for generating small neutrino masses. One of them is the type-I seesaw (or commonly referred as canonical seesaw [4]) which relies on the existence of right-handed (RH) neutrinos. Despite the success of seesaw mechanisms in explaining light neutrino masses, they suffer from direct testability at LHC or at any other near future accelerator experiments. On the other hand, one can lower the scale of new physics by generating neutrino masses through radiative mechanisms [5].

The recently observed 3.5 keV X-ray line signal in the spectrum of 73 galaxy clusters as reported by the XMM-Newton X-ray observatory [6, 7] will provide a strong hint for non-gravitational Dark Matter interaction if confirmed by others. A few attempts have been already taken in order to explain this excess of X-ray signal by (i) decaying dark matter or annihilating dark matter, [8–20], (ii) magnetic dipolar dark matter (via upscattering of dark matter into an excited state and subsequent decay of excited states into dark matter and an X-ray photon), [21–27], (iii) others [28–52]. From the recent explored scenarios it is found that such a photon signal, if the claimed excess persists, can be explained by late decay and/or annihilation of multi-keV mass dark matter, or decay of a second lightest metastable particle to lightest stable particle with a keV mass splitting. Such a monochromatic photon signal cannot be explained within the framework of standard model (SM) of particle physics.

In this paper, we attempt to explain simultaneously the 3.5 keV X-ray line and non-zero

light neutrino masses in a minimal extension of the SM while keeping the scale of new physics at the TeV scales. We extend the SM with three right-handed neutrinos (N_1, N_2, N_3), one singlet charged scalar H^+ and an extra Higgs doublet Σ . We also impose a discrete symmetry $Z_2 \times Z'_2$ which does not allow to mix the RH-neutrinos. A small mixing between N_1 and N_2 is obtained by breaking the discrete symmetry softly. As a result a small mass splitting between N_1 and N_2 is created. This allows the next-to-lightest stable particle N_2 to decay to lightest stable particle N_1 through electromagnetic dipole moment operators. If the mass splitting between N_1 and N_2 is about 3.5 keV, then the emitted photon can be identified with the observed X-ray line by XMM-Newton X-ray observatory. Since the lepton number is violated by the Majorana mass of heavy RH neutrinos, the light neutrinos acquire their masses either at tree level or at loop level.

The paper is organised as follows. In section-II and section-III we discuss in details a model of dipolar dark matter which not only give neutrino mass but also explain the 3.5 keV X-ray line. In section-IV, we discuss the relic abundance of dark matter. Constraints from direct detection of dark matter is given in section-V. Section-VI concludes.

II. THE MODEL FOR DIPOLAR DARK MATTER

We augment the SM by introducing three right-handed fermions N_{iR} which are essentially singlets under $SU(2)_L$. We also add a singlet charged scalar (H^+) and a Higgs doublet (Σ) to achieve the proposed objective. The masses of all these new particles are assumed to be of $\mathcal{O}(\text{TeV})$. A $Z_2 \times Z'_2$ symmetry is also imposed in order to keep the lightest of right handed neutrinos stable, which also serves as a candidate of dark matter. The entire particle content, along with the quantum number assignments, is displayed in Table I. We can then write the Lagrangian as:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{New}}$$

where \mathcal{L}_{SM} is the SM Lagrangian while the new physics Lagrangian \mathcal{L}_{New} contains all terms containing any of the new particles including the right handed neutrinos and is given by:

$$\begin{aligned} \mathcal{L}_{\text{New}} \ni & (Y_H)_{1\alpha} N_{1R}^T C \ell_{\alpha R} H^+ + (Y_\Sigma)_{\alpha 2} \overline{\ell_{\alpha L}} \tilde{\Sigma} N_{2R} \\ & + (Y_\nu)_{\alpha 3} \overline{\ell_{\alpha L}} \tilde{\Phi} N_{3R} + (Y_e)_{\alpha\beta} \overline{\ell_{\alpha L}} \Phi \ell_{\beta R} \\ & + \frac{1}{2} \overline{(N_{iR})^C} M_{N_i} N_{iR} + \text{h.c.} + V(\Phi, \Sigma, H^+) \end{aligned} \quad (1)$$

TABLE I: Particle content of the proposed Model.

	Field	$SU(3)_C \times SU(2)_L \times U(1)_Y$	$Z_2 \times Z'_2$
Fermions	$Q_L \equiv (u, d)_L^T$	$(3, 2, 1/6)$	$++$
	u_R	$(3, 1, 2/3)$	$++$
	d_R	$(3, 1, -1/3)$	$++$
	$\ell_L \equiv (\nu, e)_L^T$	$(1, 2, -1/2)$	$++$
	e_R	$(1, 1, -1)$	$++$
	N_{1R}	$(1, 1, 0)$	$-+$
	N_{2R}	$(1, 1, 0)$	$+ -$
	N_{3R}	$(1, 1, 0)$	$++$
Scalars	Φ	$(1, 2, +1/2)$	$++$
	Σ	$(1, 2, +1/2)$	$+ -$
	H^+	$(1, 1, +1)$	$- +$

where the scalar potential can be given by:

$$\begin{aligned}
 V(\Phi, \Sigma, H^+) &= -\mu_\Phi^2 |\Phi|^2 + M_\Sigma^2 |\Sigma|^2 + M_H^2 |H^\pm|^2 \\
 &+ \lambda_\phi |\Phi|^4 + \lambda_\Sigma |\Sigma|^4 + \lambda_H |H^\pm|^4 \\
 &+ \lambda_{\Phi H} (\Phi^\dagger \Phi) |H^\pm|^2 + \lambda_{\Sigma H} (\Sigma^\dagger \Sigma) |H^\pm|^2 \\
 &+ f |\Phi|^2 |\Sigma|^2 + \frac{\lambda_{\Phi \Sigma}}{2} [(\Phi^\dagger \Sigma)^2 + h.c.] .
 \end{aligned} \tag{2}$$

Note that the Σ is ascribed a positive mass-squared so that it does not acquire any vacuum expectation value (vev). Furthermore, we assume that $M_\Sigma > M_2$ and $M_{H^\pm} > M_1$ so that the decays $N_1 \rightarrow \ell_R H^+$ and $N_2 \rightarrow \nu_L + \Sigma^0$ are kinematically forbidden. As a result N_1 and N_2 are individually stable. However, under $Z_2 \times Z'_2$ symmetry, N_3 goes to itself and hence it is not stable. It can decay through the process: $N_3 \rightarrow \nu_i \Phi^0$. After Electro-Weak phase transition, the neutrino can acquire a small Majorana mass through the type-I seesaw anchored by N_3 .

Notice that $Z_2 \times Z'_2$ symmetry doesn't allow N_1 and N_2 to mix with each other. We generate a small mixing between them by breaking $Z_2 \times Z'_2$ softly with:

$$\mathcal{L}_{\text{soft}} = [\mu_s \Sigma \Phi (H^+)^* + h.c.] , \tag{3}$$

where μ_s has to be determined from the observed phenomenon. In particular, the breaking of $Z_2 \times Z'_2$ symmetry allows a mixing between N_1 and N_2 . As a result N_2 can decay via $N_2 \rightarrow N_1 + \gamma$. The emitted photon can be identified with the recent observation of 3.5 keV X-ray line signal by the XMM-Newton X-ray observatory.

A. Constraints on new particles

Since Σ is a scalar doublet under $SU(2)_L$, it couples to Z -boson and hence can modify the decay width of Z -boson. Therefore, we take the masses of Σ particles to be larger than $M_Z/2$. That means

$$M_\Sigma > 45\text{GeV} . \quad (4)$$

The mass of the singlet charged scalar H^\pm is lower bounded by LEP. It is given to be [53]

$$M_H > 80\text{GeV} . \quad (5)$$

B. Mass splitting between N_1 and N_2




FIG. 1: Mass splitting between N_1 and N_2 due to radiative correction through breaking of $Z_2 \times Z'_2$ symmetry.

As mentioned before, the breaking of $Z_2 \times Z'_2$ symmetry generates a mixing between N_1 and N_2 . The mixing can be calculated from Fig. 1 as:

$$\Delta M_{12} = \frac{Y_{H1\alpha}^* Y_{\Sigma\alpha 2}}{16\pi^2} \frac{\mu_s v_{ew} m_\ell}{(M_\Sigma^2 - M_H^2)} \ln \left(\frac{M_\Sigma^2}{M_H^2} \right) \quad (6)$$

As a result the Majorana mass matrix in the basis of (N_1, N_2, N_3) can be given by:

$$\begin{pmatrix} M_1 & \Delta M_{12} & 0 \\ \Delta M_{12} & M_2 & 0 \\ 0 & 0 & M_3 \end{pmatrix} \quad (7)$$

Diagonalizing the mass matrix, we get the mass eigenvalues $M_1 + \Delta M_{12}$ and $M_2 - \Delta M_{12}$ and M_3 . Thus the mass splitting between the two eigenstates N_1 and N_2 is given by

$$\delta = 2\Delta M_{12}. \quad (8)$$

If we assume that N_2 is heavier than N_1 , then the former can decay to latter by emitting a monochromatic photon with energy equal to the mass difference between them. Moreover, the life time of N_2 should be larger than the age of the universe so that it can decay in the current epoch. If the mass difference between N_2 and N_1 is of \sim keV, then the recent observation of X-ray line can be accommodated in this beyond SM scenario.

III. MAGNETIC DARK MATTER (MDM) AND MONOCHROMATIC PHOTON

Let us first discuss the electromagnetic properties of heavy RH neutrinos N_i with photon before deducing the potential implications of the magnetic dipole moment operator in radiative decay $N_2 \rightarrow N_1 + \gamma$. Due to the Majorana nature, the diagonal magnetic moment of heavy Majorana neutrinos is zero. There is only transition magnetic moment for them. The electromagnetic coupling for the heavy neutrinos with photon, via dimension-five effective magnetic dipole moment (MDM) operator is

$$\mathcal{L}_{\text{MDM}} = -\frac{i}{2} N_{Rj} C^{-1} \mu_{jk} \sigma_{\alpha\beta} N_{Rk} F^{\alpha\beta} + \text{h.c.} \quad (j \neq k). \quad (9)$$

where $\mathcal{F}^{\alpha\beta}$ is the electromagnetic field tensor and μ_{12} is the transition magnetic moment between the first and second generation of heavy right-handed Majorana neutrinos N_2, N_1 which can be calculated from Fig. (2) as

$$\mu_{12} = -\frac{-e}{64 \pi^2} \frac{(Y_H^* Y_\Sigma \mu_s v_{\text{ew}})}{M_\Sigma^2 - M_H^2} \frac{m_\ell}{M_\Sigma^2} \mathcal{I}_{\text{tot}} \quad (10)$$




FIG. 2: Diagram contributing to transient magnetic moment of right-handed neutrino in the proposed model.

where

$$\mathcal{I}_{\text{tot}} \simeq \int_0^1 dx \left[\frac{x(1-x)^2}{(1-x)M_H^2/M_\Sigma^2 + x(x-1)M_2^2/M_\Sigma^2 + x} - \frac{x(1-x)^2}{(1-x) + x(x-1)M_2^2/M_\Sigma^2 + xM_H^2/M_\Sigma^2} \right]. \quad (11)$$

Assuming $M_2 < M_\Sigma, M_H$ and with a typical set of values: $M_2 \simeq 100$ GeV, $M_\Sigma \simeq M_H = 384$ GeV, the loop factor is estimated to be 4.4×10^{-5} .

Since we assume the mass splitting between N_1 and N_2 is order of keV, the decay of N_2 to N_1 can give rise a monochromatic X -ray line. The decay rate of $N_2 \rightarrow N_1 \gamma$ is estimated as:

$$\begin{aligned} \Gamma(N_2 \rightarrow N_1 \gamma) &= \frac{M_2^3}{8\pi} \left(1 - \frac{M_1^2}{M_2^2}\right)^3 |\mu_{12}|^2 \\ &= \frac{|\mu_{12}|^2}{\pi} \delta^3, \end{aligned} \quad (12)$$

where

$$\delta \equiv E_\gamma = \frac{M_2}{2} \left(1 - \frac{M_1^2}{M_2^2}\right) \quad (13)$$

is the energy of the emitted photon, which is nothing but the mass difference between N_1 and N_2 . For the observed X -ray line we use $\delta \sim 3.5$ keV and $\Gamma(N_2 \rightarrow N_1 \gamma) = 0.36 - 3.3 \times 10^{-52} \text{ GeV} (M_2/3.5 \text{ keV})$ for decaying dark matter [6, 13, 27]. This implies that for $M_2 = 100$ GeV, the decay rate is $\mathcal{O}(10^{-44}) \text{ GeV}$. In other words the life time of N_2 is $\mathcal{O}(10^{19}) \text{ sec}$,

which is larger than the age of the Universe. From Eqn. (12) one can estimate the required magnetic moment is $\mu_{12} = \mathcal{O}(10^{-14}) \text{ GeV}^{-1}$.

A. Scale of new physics and collider search

In order to fix the scale of new physics let us define the ratio:

$$R \equiv \frac{\mu_{12}}{\delta} = \frac{e}{8} \frac{1}{M_\Sigma^2} \frac{I_{\text{tot}}}{\ln(M_\Sigma^2/M_H^2)}. \quad (14)$$

In the above equation I_{tot} can be evaluated numerically. In particular, for a 350 GeV DM mass, using $\delta = 3.5 \text{ keV}$ and $\mu_{12} = 2.46 \times 10^{-14} \text{ GeV}^{-1}$ we get $M_\Sigma \approx M_H = 380 \text{ GeV}$. Thus the mass scale of the new particles are not far from the electroweak scale and hence can be searched at the collider. In particular, the charged scalars H^\pm and Σ^\pm are important. These particles can be pair produced at LHC via the exchange of SM Higgs particle. For example, $pp \rightarrow h \rightarrow H^+H^- \rightarrow e^+e^- + \text{missing energy}$. Similarly, $pp \rightarrow h \rightarrow \Sigma^+\Sigma^- \rightarrow e^+e^- + \text{missing energy}$. Σ^\pm particles can also be detected through other decay processes, such as: $\Sigma^\pm \rightarrow W^\pm \Sigma^0 (\Sigma^{0*}) \rightarrow f \bar{f} f_1 \bar{f}_2$, where f, f_1, f_2 are SM fermions.

B. Light neutrino Mass

Through the electroweak (EW) phase transition, one of the Yukawa term $(Y_\nu)_{\alpha 3} \bar{\ell}_{\alpha L} \tilde{\Phi} N_{3R}$ generates a Dirac mass term: $(M_D)_{\alpha 3} = (Y_\nu)_{\alpha 3} v_{\text{ew}}$, where $v_{\text{ew}} = \langle \Phi \rangle$ and $\alpha = e, \mu, \tau$. But the N_3 has a Majorana mass term $M_3 N_3 N_3$ which breaks lepton number by two units. As a result we get a Majorana mass matrix for light neutrinos to be:

$$(m_\nu)_{\alpha\beta} = \frac{v^2}{M_3} (Y_\nu)_{\alpha 3} (Y_\nu)_{\beta 3} \quad (15)$$

Diagonalizing the above mass matrix we get the eigenvalues: $\text{Tr} [(Y_\nu)_{\alpha 3} (Y_\nu)_{\beta 3}] v_{\text{ew}}^2 / M_3, 0, 0$. Thus at the tree level only one of the neutrinos is massive, say ν_3 . This is because of the exact $Z_2 \times Z_2'$ symmetry in the Lagrangian which prevents the Yukawa terms $\bar{\ell}_\alpha \Phi N_{1R}$ and $\bar{\ell}_\alpha \Phi N_{2R}$. On the other hand, the fact that $M_\Sigma^2 > 0$ prevents a vacuum expectation value for Σ field, we can not generate a Majorana mass of light neutrinos at the tree level through the coupling: $Y_{\Sigma \alpha 2} \bar{\ell}_{\alpha L} \tilde{\Sigma} N_{2R}$. However, the latter can be generated through one loop radiative correction diagram as shown in Fig.3. Because of the quartic coupling: $\frac{\lambda_{\Phi\Sigma}}{2} [(\Phi^\dagger \Sigma)^2 + h.c.]$,




FIG. 3: One loop diagram contributing to Majorana mass of light neutrinos.

the EW phase transition generates a mass square splitting: $\lambda_{\Phi\Sigma}v^2$ between the real (Σ_r^0) and imaginary (Σ_I^0) components of Σ^0 , the neutral component of Σ field. As a result, the analytical expression for one loop-generated light neutrino mass is given by [54]:

$$(m_{\nu}^{\text{loop}})_{\alpha\beta} = \frac{(Y_{\Sigma})_{\alpha 2}(Y_{\Sigma})_{\beta 2} M_2}{16\pi^2} \left[\frac{M_{\Sigma_R}^2}{(M_{\Sigma_R}^2 - M_2^2)} \ln \left(\frac{M_{\Sigma_R}^2}{M_2^2} \right) - \frac{M_{\Sigma_I}^2}{(M_{\Sigma_I}^2 - M_2^2)} \ln \left(\frac{M_{\Sigma_I}^2}{M_2^2} \right) \right] \quad (16)$$

Diagonalising the above radiative mass matrix we get only one of the state massive, say ν_2 . Since the origin of the masses of the two eigen states ν_2 and ν_3 are different they can easily satisfy the solar and atmospheric mass splitting constraint.

IV. RELIC ABUNDANCE OF N_1 AND N_2

The lightest stable particle (LSP), which is odd under $Z_2 \times Z'_2$ symmetry, is the N_1 and hence behaves as a candidate of dark matter. The next to lightest stable particle (NLSP) is the N_2 and is stable on the cosmological time scale. It decays to N_1 through the electromagnetic coupling as shown in Fig. 2. Since the observed X-ray line implies that the effective electromagnetic coupling of N_1 and N_2 with photon is extremely small, the life time of N_2 is larger than the age of the Universe. Therefore, N_2 is also a candidate of dark matter. When N_2 decays its density gets converted to N_1 , while the net DM abundance remain intact.

Since the electromagnetic coupling between N_1 , N_2 and γ is small, it can not bring the former two to thermal equilibrium. Note that to bring N_1 and N_2 to thermal equilibrium through electromagnetic coupling one needs the magnetic dipole moment $\mu_{12} \sim 10^{-4}\text{GeV}^{-1}$ [55]. Hence the only way the right handed neutrinos are brought to thermal

equilibrium is the interaction of N_1 with the singlet charged Higgs (H^+) and right-handed charged leptons (ℓ_R) and that of N_2 with the $SU(2)_L$ doublets ℓ_L and Σ . For estimating the relic abundance of DM, the keV scale mass splitting between N_1 and N_2 is irrelevant. So for all practical purpose we assume $M_{N_1} = M_{N_2} = M_{\text{DM}}$. As the temperature falls below the mass of DM, it decouples from the thermal bath. The decoupling temperature is given by $T_f = M_{\text{DM}}/x_f$, where $x_f \approx 25$. The relevant cross-sections are given in appendix-B. The relic abundance obtained through co-annihilation of N_1 with H^+ and that of N_2 with Σ can be given by [56]

$$\Omega_{\text{DM}} h^2 = \frac{1.09 \times 10^9 \text{GeV}^{-1}}{g_*^{1/2} M_{\text{Pl}}} \times \frac{1}{J(x_f)} \quad (17)$$

where

$$J(x_f) = \int_{x_f}^{\infty} \frac{\langle \sigma|v| \rangle_{\text{eff}}}{x^2} \quad (18)$$

with

$$\langle \sigma|v| \rangle_{\text{eff}} = \langle \sigma|v| \rangle_{\text{eff}}^{N_1} + \langle \sigma|v| \rangle_{\text{eff}}^{N_2}. \quad (19)$$

In the above equation the effective co-annihilation cross-sections of N_1 with H^+ and that of N_2 with that of Σ are given as:

$$\langle \sigma|v| \rangle_{\text{eff}}^{N_i} = \langle \sigma|v| \rangle^{N_i-X} g_i g_X (1 + \Delta_X)^{3/2} \times \left(\frac{e^{-x\Delta_X}}{g_{\text{eff}}^2} \right) \quad (20)$$

where $g_i, i = 1, 2$ represents the internal degrees of freedom of N_1 and N_2 , g_X represents the internal degrees of freedom of X -particle (H^+, Σ) co-annihilating with $N_i, i = 1, 2$ and

$$g_{\text{eff}} = g_i + g_X (1 + \Delta_X)^{3/2} e^{-x\Delta_X}. \quad (21)$$

where $\Delta_X = (M_X - M_{\text{DM}})/M_{\text{DM}}$.

To estimate the contribution of N_1 and N_2 to relic abundance we define the ratios:

$$\begin{aligned} r_1 &= \frac{\langle \sigma|v| \rangle_{N_1}}{\langle \sigma|v| \rangle_{N_1} + \langle \sigma|v| \rangle_{N_2}} \\ r_2 &= \frac{\langle \sigma|v| \rangle_{N_2}}{\langle \sigma|v| \rangle_{N_1} + \langle \sigma|v| \rangle_{N_2}} \end{aligned} \quad (22)$$

We have shown r_1 and r_2 as a function of M_{DM} in Fig. (4) for $Y_{\Sigma} = Y_H = Y$. Moreover, for simplicity we assume the masses of H^+ and Σ to be M_X . We define $\Omega_{\text{old}} h^2$ as the relic abundance of DM assuming $M_{\text{DM}} = M_X$. Then we check the effect of mass splitting on the relic abundance. This is shown in Fig. (5). From Fig. (5) we see that relic abundance




FIG. 4: r_1 and r_2 as a function of M_{DM} for $Y_{\Sigma} = Y_H$. For simplicity we set $M_{\text{DM}} = M_X$.

increases with the rise of mass splitting between the DM and X-particles. This is due to the fact that relic abundance is inversely proportional to the effective cross-section, which suffers a Boltzmann suppression, i.e., $\langle\sigma|v|\rangle \propto e^{-x\Delta X}$. Therefore, the effective cross-section decreases with the increase of mass splitting. Next we show the dependency of the observed




FIG. 5: Ratio of dark matter abundance as a function of $M_X - M_{\text{DM}}$ for a typical mass of DM, $M_{\text{DM}} = 1000\text{GeV}$ and $Y_{\Sigma} = Y_H = 0.55$

relic abundance simultaneously on the mass of DM and the mass splitting between DM and X-particles for various couplings. This is demonstrated in Fig. (6) for $Y = 0.4, 0.5, 0.6$. As expected for small mass splitting the co-annihilation is large between N_1, N_2 and the corresponding X-particles. Therefore we get large number of points near $M_X - M_{\text{DM}} \approx 0$. On the other hand, for a given mass of DM, if the coupling increases then we need relatively large mass splitting so that the effective cross-section decreases to get the right relic abundance. This feature can be easily read from Fig. (6) as we move from $Y = 0.4$ to $Y = 0.6$. For light mass of DM, say $M_{\text{DM}} < 100\text{GeV}$, the cross-section increases with the further decrease

of DM mass. Therefore, we need a large mass splitting so that the effective cross-section decreases to give the correct relic abundance.




FIG. 6: Observed relic abundance in the plane of $M_X - M_{DM}$ versus M_{DM} for different values of the couplings, i.e., $Y = 0.4, 0.5, 0.6$.

V. DIRECT DETECTION OF DARK MATTER AND CONSTRAINTS

In this section, we wish to constrain the model parameters from the direct detection experiments for dark matter and LUX experiment [57] is one such experiment considered in our present analysis. Since the mass splitting between N_1 and N_2 is only 3.5 keV, so for all practical purposes we consider the spin independent DM-nucleon interaction $N_1 n \rightarrow N_2 n$, mediated by SM Higgs exchange, to be elastic. The Feynman diagram for DM-nucleon interaction for direct detection is shown in Fig. 7. We denote the effective coupling between $N_2 - N_1 - \Phi$ entering into this interaction as λ_{eff} and is given by:

$$\lambda_{\text{eff}} \simeq \frac{-\mu_S}{16\pi^2} Y_\Sigma Y_H^* m_\ell \mathcal{F}(M_H^2, M_\Sigma^2, m_\ell^2), \quad (23)$$

where

$$\mathcal{F}(M_H^2, M_\Sigma^2, m_\ell^2) = \frac{1}{M_H^2 - M_\Sigma^2} \times \left[\frac{1}{\frac{m_\ell^2}{M_H^2} - 1} \ln\left(\frac{m_\ell^2}{M_H^2}\right) - \frac{1}{\frac{m_\ell^2}{M_\Sigma^2} - 1} \ln\left(\frac{m_\ell^2}{M_\Sigma^2}\right) \right]. \quad (24)$$

Thus the spin independent DM-nucleon cross-section can be given as:

$$\sigma_{SI} = \frac{\mu_n^2}{4\pi} \lambda_{\text{eff}}^2 \left(\frac{f_n m_n}{v_{\text{ew}}} \right)^2 \frac{1}{M_h^4} \quad (25)$$




FIG. 7: Elastic scattering of magnetic dipolar dark matter with target nuclei.

where $\mu_n = M_{N_1} m_n / (M_{N_1} + m_n)$ is the reduced DM-nucleon mass with $m_n = 0.946 \text{ GeV}$ and $M_h = 125 \text{ GeV}$ is the SM Higgs mass. The effective coupling between the SM Higgs and nucleon is given by: $f_n m_n / v_{ew}$ which depend upon the quark content of the nucleon for each quark flavour. The Higgs-nucleon coupling f_n is given by:

$$f_n = \sum_q f_q = \frac{m_q}{m_n} \langle n | \bar{q}q | n \rangle \quad (26)$$

where the sum is over all quark flavors. In the present analysis, we have used $f_n = 0.32$ [58] though its value can lie within a range $f_n = 0.26 - 0.33$ [59]. Using $M_\Sigma \approx M_H = 100 \text{ GeV}$ and $Y_H = Y_\Sigma = 0.5$ we get $\sigma_{\text{SI}} \approx \mathcal{O}(10^{-59}) \text{ cm}^2$, which is much smaller than the current stringent limit: $\sigma_{\text{SI}} = 7.5 \times 10^{-46} \text{ cm}^2$ from LUX.

VI. CONCLUSION

In this paper we have shown that the observed 3.5 keV X -ray line signal and neutrino mass can be explained simultaneously in a minimal extension of the SM. We extend the SM with three right-handed neutrinos (N_1, N_2, N_3), a charged scalar H^+ and a Higgs doublet (Σ) which transforms non-trivially under a discrete symmetry $Z_2 \times Z'_2$. The lightest odd particle is N_1 and behaves as a candidate of DM. The next to lightest odd particle is N_2 whose mass is separated from N_1 by 3.5 keV. As a result the total DM of the universe is a mixture of N_1 and N_2 abundance. However, the discrete symmetry is broken softly so that N_2 decays to $N_1 + \gamma$ in the present universe, converting its abundance to N_1 while keeping the total DM abundance intact. The γ can be identified with the observed 3.5 keV X -ray line. Since N_3 couples to the SM lepton and Higgs doublet, a small Majorana mass for one of the light

neutrinos is generated through the canonical seesaw mechanism. The other neutrinos get their masses at one loop level.

We show that the required relic abundance of DM implies the masses of new particles H^\pm and Σ^\pm are not far from the electroweak scale. So these particles can be pair produced and their subsequent decay can be searched at LHC.

We noticed that the direct detection cross-section of DM with the nucleon is highly suppressed. Therefore, the constraints on the model parameters are almost negligible.

VII. ACKNOWLEDGMENTS

The work of Sudhanwa Patra is partially supported by the Department of Science and Technology, Govt. of India under the financial grant SB/S2/HEP-011/2013. Narendra Sahu is partially supported by the Department of Science and Technology, Govt. of India under the financial Grant SR/FTP/PS-209/2011.

Appendix A: Calculation for transitional magnetic moment for magnetic dark matter decay $N_2 \rightarrow N_1 \gamma$

The effective Lagrangian accounting the interaction between photon and two neutral fermions states is given as

$$\mathcal{L}_{\text{MDM}} = -\frac{i}{2} N_{Rj} C^{-1} \mu_{jk} \sigma_{\alpha\beta} N_{Rk} F^{\alpha\beta} + \text{h.c.} \quad (j \neq k). \quad (27)$$

We present here the Feynman calculation for transitional magnetic moment between two neutral fermion states N_2 and N_1 leading to the decay $N_2 \rightarrow N_1 \gamma$ explaining the 3.5 keV X-ray line signal. The possible Feynman diagrams which yield transitional magnetic moment μ_{12} and decay process $N_2 \rightarrow N_1 \gamma$ is shown in Fig. 8. Let us assign the four-vector momenta of N_2 , N_1 and photon by p_1 , p_2 and k , respectively. The Feynman amplitude for the magnetic dark matter decay shown in Fig. 8(a) can be written as

$$\begin{aligned} \mathcal{M}^{(a)} &= \int \frac{d^4 k}{(2\pi)^4} \overline{u(p_2)} (i Y_H^* P_R) S_\ell(k - q) (-i e \gamma_\mu \epsilon^\mu(q)) S_\ell(k) (i Y_\Sigma P_R) u^c(p_1) \\ &\quad \times \Delta_H(p_1 - k) (\mu_s v_{\text{ew}}) \Delta_\Sigma(p_1 - k) \\ &= \overline{u(p_2)} (i \Gamma_\mu^{(a)}) \epsilon^\mu(q) u^c(p_1), \end{aligned} \quad (28)$$




FIG. 8: Feynman diagram for magnetic dark matter decay: $N_2 \rightarrow N_1 \gamma$.

where $i \Gamma_\mu^{(a)}$ is factored out to be

$$\begin{aligned}
 i \Gamma_\mu^{(a)} &= (-e Y_H^* Y_\Sigma \mu_s v_{ew}) \times \\
 &\int \frac{d^4 k}{(2\pi)^4} \frac{P_R (\not{k} - \not{q} + m_\ell) \gamma_\mu (\not{k} + m_\ell) P_R}{[\{(k-q)^2 - m_\ell^2\} \{k^2 - m_\ell^2\} \{(p_1 - k)^2 - M_H^2\} \{(p_1 - k)^2 - M_\Sigma^2\}]} \\
 &= (-e Y_H^* Y_\Sigma \mu_s v_{ew}) \int \frac{d^4 k}{(2\pi)^4} \mathcal{D}^{-1} \cdot \mathcal{N}_\mu
 \end{aligned} \tag{29}$$

We denote \mathcal{D}^{-1} and \mathcal{N}_μ as follows

$$\mathcal{D}^{-1} = \frac{1}{[\{(k-q)^2 - m_\ell^2\} \{k^2 - m_\ell^2\} \{(p_1 - k)^2 - M_H^2\} \{(p_1 - k)^2 - M_\Sigma^2\}]} \tag{30}$$

$$\mathcal{N}_\mu = P_R (\not{k} - \not{q} + m_\ell) \gamma_\mu (\not{k} + m_\ell) P_R. \tag{31}$$

After doing some simpler algebra, we get

$$\begin{aligned}
 \Gamma_\mu^{(a)} &= \frac{(-e Y_H^* Y_\Sigma \mu_s v_{ew})}{M_\Sigma^2 - M_H^2} \times \int \frac{d^4 k}{(2\pi)^4} \left[\frac{N_\mu}{[\{(k-q)^2 - m_\ell^2\} \{k^2 - m_\ell^2\} \{(p_1 - k)^2 - M_\Sigma^2\}]} \right. \\
 &\quad \left. - \frac{\mathcal{N}_\mu}{[\{(k-q)^2 - m_\ell^2\} \{k^2 - m_\ell^2\} \{(p_1 - k)^2 - M_H^2\}]} \right] \\
 &= \frac{(-e Y_H^* Y_\Sigma \mu_s v_{ew})}{M_\Sigma^2 - M_H^2} [I_1 - I_2]
 \end{aligned} \tag{32}$$

The integral now becomes

$$\begin{aligned}
 I_1 &= \frac{m_\ell}{16 \pi^2} \int_0^1 dx \int_0^{1-x} dy S_1^{-2} [2(qx + p_1 y)_\mu - \not{q} \gamma_\mu] P_R \\
 &= \frac{1}{16 \pi^2} \frac{m_\ell}{M_\Sigma^2} \int_0^1 dx \int_0^{1-x} dy \Omega_1 [2y p_{1\mu} - 2p_{1\mu}] P_R \\
 &= \frac{1}{8 \pi^2} \frac{m_\ell}{M_\Sigma^2} p_{1\mu} \int_0^1 dx \int_0^{1-x} dy \Omega_1 [y - 1] P_R
 \end{aligned}$$

Here $S_1^{-2} = M_\Sigma^{-2} \Omega_1$, $\Omega_1 = [Z_N (y - y^2 - xy) - (1 - y)Z_\ell - y]^{-1}$. Since the term proportional to $p_{2\mu} - p_{1\mu}$ gives vanishing contribution and writing $p_{1\mu} = \frac{1}{2}[p_{2\mu} + p_{1\mu} + q_\mu]$, the integral becomes

$$I_1 = \frac{1}{16\pi^2} \frac{m_\ell}{M_\Sigma^2} (p_1 + p_2)_\mu \int_0^1 dx \int_0^{1-x} dy \Omega_1 [y - 1] P_R \quad (33)$$

Similarly, one can express the second integral \mathcal{I}_2 by replacing M_Σ by M_H .

Now we can write the loop factor as

$$\begin{aligned} i\Gamma_\mu^{(a)} &= \frac{(-e Y_H^* Y_\Sigma \mu_s v_{ew})}{M_\Sigma^2 - M_H^2} \left[\frac{1}{16\pi^2} \frac{m_\ell}{M_\Sigma^2} (p_1 + p_2)_\mu \int_0^1 dx \int_0^{1-x} dy \Omega_1 (y - 1) P_R \right. \\ &\quad \left. - \frac{1}{16\pi^2} \frac{m_\ell}{M_\Sigma^2} (p_1 + p_2)_\mu \int_0^1 dx \int_0^{1-x} dy \Omega_2 (y - 1) P_R \right] \\ &= \frac{-e (Y_H^* Y_\Sigma \mu_s v_{ew})}{16\pi^2} \frac{m_\ell}{M_\Sigma^2 - M_H^2} \frac{m_\ell}{M_\Sigma^2} \mathcal{I} (p_1 + p_2)_\mu P_R \\ &= \mathcal{A}_{12}^{(a)} (p_1 + p_2)_\mu \end{aligned} \quad (34)$$

where the integral \mathcal{I} is

$$\mathcal{I} = \int_0^1 dx \int_0^{1-x} dy (y - 1) [\Omega_1 - \Omega_2]$$

and

$$\mathcal{A}_{12}^{(a)} = \frac{-e (Y_H^* Y_\Sigma \mu_s v_{ew})}{16\pi^2} \frac{m_\ell}{M_\Sigma^2 - M_H^2} \frac{m_\ell}{M_\Sigma^2} \mathcal{I}$$

From Gordan Identity, we can write

$$\mathcal{A}_{12}^{(a)} \overline{u(p_2)} (p_1 + p_2)_\mu u(p_1) = \mathcal{A}_{12}^{(a)} \overline{u(p_2)} [2M_N \gamma_\mu - i\sigma_{\mu\nu} (p_2 - p_1)^\nu] u(p_1) \quad (35)$$

The analytical expression for transitional magnetic moment between two nearly degenerate heavy RH neutrinos, as derived from the effective operator $\mathcal{A}_{12}^{(a)} \overline{N_1} \sigma^{\mu\nu} N_2 F_{\mu\nu}$ following Feynman diagram Fig. 8(a), as

$$\mathcal{A}_{12}^{(a)} = \frac{-e (Y_H^* Y_\Sigma \mu_s v_{ew})}{16\pi^2} \frac{m_\ell}{M_\Sigma^2 - M_H^2} \frac{m_\ell}{M_\Sigma^2} \mathcal{I} \quad (36)$$

Following the same method of Feynman calculation, one can derive the relevant contributions of transition magnetic moment depicted in Fig. 8(b). At the end, the analytical expression for transitional magnetic moment including relevant Feynman diagrams can be put in the following expression

$$\begin{aligned} \mu_{12} &= \mathcal{A}_{12}^{(a)} + \mathcal{A}_{12}^{(b)} \\ &\simeq \frac{-e (Y_H^* Y_\Sigma \mu_s v_{ew})}{16\pi^2} \frac{m_\ell}{M_\Sigma^2 - M_H^2} \frac{m_\ell}{M_\Sigma^2} \mathcal{I}_{\text{tot}}. \end{aligned} \quad (37)$$

Appendix B: Cross-section calculation for relic abundance of DM

We present here the cross-section calculation involving magnetic dark matter and possible coannihilation channels relevant for relic abundance estimation. The relic abundance of N_1 is obtained through its co-annihilation with H^+ via the following processes:

$$\langle\sigma|v\rangle(N_{1R} + H^+ \Leftrightarrow \ell_R + \phi) = \frac{(Y_H)_{1\alpha}^2 m_f^2}{64\pi M_H (s - m_\alpha^2)^2 \langle\phi\rangle^2} s\sqrt{s} \left(1 - \frac{M_\phi^2}{s}\right)^2 \quad (38)$$

$$\langle\sigma|v\rangle(N_{1R} + H^+ \Leftrightarrow \ell_R + Z) = \frac{1}{32\pi M_H} \left(\frac{g(Y_H)_{1\alpha}(T_3 - \sin^2\theta_W Q)}{\cos\theta_W (s - m_\alpha^2 + i\epsilon)}\right)^2 s\sqrt{s} \left(1 - \frac{m_Z^2}{s}\right)^2 \quad (39)$$

$$\langle\sigma|v\rangle(N_{1R} + H^+ \Leftrightarrow \ell_R + \gamma) = \frac{e^2}{32\pi M_H} \frac{(Y_H)_{1\alpha}^2}{(s - m_\alpha^2)^2} s\sqrt{s} \quad (40)$$

where $\alpha = e, \mu, \tau$

The relic abundance of N_2 is obtained through its co-annihilation with Σ particles via the following processes are

$$\langle\sigma|v\rangle(N_{2R} + \Sigma^0 \Leftrightarrow l_L + W) = \frac{1}{64\pi M_\Sigma} \frac{g^2(Y_\Sigma)_{2\alpha}^2}{\sqrt{s}} \left(1 - \frac{m_W^2}{s}\right)^2 \quad (41)$$

$$\langle\sigma|v\rangle(N_{2R} + \Sigma^0 \Leftrightarrow \nu_L + Z) = \frac{1}{128\pi M_\Sigma} \frac{g^2(Y_\Sigma)_{2\alpha}^2}{\cos^2\theta_W \sqrt{s}} \left(1 - \frac{m_Z^2}{s}\right)^2 \quad (42)$$

$$\langle\sigma|v\rangle(N_{2R} + \Sigma^- \Leftrightarrow \nu_L + W) = \frac{1}{64\pi M_\Sigma} \frac{g^2(Y_\Sigma)_{2\alpha}^2}{(s - m_\alpha^2 + i\epsilon)^2} s\sqrt{s} \left(1 - \frac{m_W^2}{s}\right)^2 \quad (43)$$

$$\langle\sigma|v\rangle(N_{2R} + \Sigma^- \Leftrightarrow e_L + Z) = \frac{1}{32\pi M_\Sigma} \frac{g^2(Y_\Sigma)_{2\alpha}^2 (T_3 - \sin^2\theta_W Q)^2}{\cos^2\theta_W (s - m_\alpha^2 + i\epsilon)^2} s\sqrt{s} \left(1 - \frac{m_Z^2}{s}\right)^2 \quad (44)$$

$$\langle\sigma|v\rangle(N_{2R} + \Sigma^- \Leftrightarrow e_L + \gamma) = \frac{e^2(Y_\Sigma)_{2\alpha}^2}{32\pi M_\Sigma (s - m_\alpha^2 + i\epsilon)^2} s\sqrt{s} \quad (45)$$

$$\langle\sigma|v\rangle(N_{2R} + \Sigma^- \Leftrightarrow e_R + \phi) = \frac{(Y_\Sigma)_{2\alpha}^2 m_f^2}{64\pi M_\Sigma (s - m_\alpha^2 + i\epsilon)^2 \langle\phi\rangle^2} s\sqrt{s} \left(1 - \frac{M_\phi^2}{s}\right)^2 \quad (46)$$

-
- [1] P. A. R. Ade *et al.* [Planck Collaboration], *Planck 2013 results. XVI. Cosmological parameters*, *Astron. Astrophys.* **571**, A16 (2014), arXiv:1303.5076 [astro-ph.CO].
- [2] G. Bertone, D. Hooper and J. Silk, *Particle Dark Matter: Evidence, Candidates and Constraints*, *Phys. Rept.* 405, 279 (2005), arXiv:hep-ph/0404175; G. Jungman, M. Kamionkowski and K. Griest, *Supersymmetric Dark Matter*, *Phys. Rept.* 267, 195 (1996), arXiv:hep-ph/9506380.

- [3] S. Fukuda et al. (Super-Kamiokande), Phys. Rev. Lett. **86**, 5656 (2001); Q. R. Ahmad et al. (SNO), Phys. Rev. Lett. **89**, 011301 (2002); Phys. Rev. Lett. **89**, 011302 (2002); J. N. Bahcall and C. Pena-Garay, New J. Phys. **6**, 63 (2004); K. Abe et al. [T2K], Phys. Rev. Lett. **107**, 041801 (2011); P. Adamson et al. [MINOS], Phys. Rev. Lett. **107**, 181802 (2011); Y. Abe et al. [DOUBLE-CHOOZ], Phys. Rev. Lett. **108**, 131801 (2012); F. P. An et al. [DAYA-BAY], Phys. Rev. Lett. **108**, 171803 (2012); J. K. Ahn et al. [RENO], Phys. Rev. Lett. **108**, 191802 (2012).
- [4] P. Minkowski, Phys. Lett. B **67**, 421 (1977); M. Gell-Mann, P. Ramond and R. Slansky in Supergravity (P. van Nieuwenhuizen and D. Freedman, eds), (Amsterdam), North Holland, 1979; T. Yanagida in Workshop on Unified Theory and Baryon number in the Universe (O. Sawada and A. Sugamoto, eds), (Japan), KEK 1979; R.N. Mohapatra and G. Senjanovic, Phys.Rev. Lett. **44**, 912 (1980).
- [5] A. Zee, Phys. Lett. B **93**, 389 (1980) [Erratum-ibid. B **95**, 461 (1980)]; K. S. Babu, Phys. Lett. B **203**, 132 (1988); E. Ma, Phys. Rev. Lett. **81**, 1171 (1998); arXiv:9805219 [hep-ph].
- [6] E. Bulbul, M. Markevitch, A. Foster, R. K. Smith, M. Loewenstein, *et al.*, Astrophys. J. **789** (2014) 13; arXiv:1402.2301 [astro-ph.CO]. A. Boyarsky, O. Ruchayskiy, D. Iakubovskiy and J. Franse, arXiv:1402.4119 [astro-ph.CO].
- [7] XMM-Newton Science Analysis System, <http://xmm.esa.int/sas/>
- [8] K.N. Abazajian, *Resonantly-Produced 7 keV Sterile Neutrino Dark Matter Models and the Properties of Milky Way Satellites*, Phys. Rev. Lett. **112**, 161303 (2014); arXiv:1403.0954 [astro-ph.CO].
- [9] S. Baek and H. Okada, *7 keV Dark Matter as X-ray Line Signal in Radiative Neutrino Model*, arXiv:1403.1710 [hep-ph].
- [10] B. Shuve and I. Yavin, *A Dark Matter Progenitor: Light Vector Boson Decay into (Sterile) Neutrinos*, Phys. Rev. **D 89**, 113004 (2014); [arXiv:1403.2727].
- [11] T. Tsuyuki, *Neutrino masses, leptogenesis, and sterile neutrino dark matter*, Phys. Rev. **D 90**, 013007 (2014); arXiv:1403.5053.
- [12] F. Bezrukov and D. Gorbunov, *Relic Gravity Waves and 7 keV Dark Matter from a GeV scale inflaton*, arXiv:1403.4638 [hep-ph].
- [13] K. P. Modak, *3.5 keV X-ray Line Signal from Decay of Right-Handed Neutrino due to Transition Magnetic Moment*, arXiv:1404.3676 [hep-ph].

- [14] D. J. Robinson and Y. Tsai, *A Dynamical Framework for KeV Dirac Neutrino Warm Dark Matter*, Phys. Rev. **D 90**, 045030 (2014); arXiv:1404.7118 [hep-ph].
- [15] S. Chakraborty, D.K. Ghosh and S. Roy, *7 keV sterile neutrino dark matter in $U(1)_R$ -lepton number model*, JHEP 10 (2014) 146; arXiv:1405.6967 [hep-ph].
- [16] A. Abada, G. Arcadi, M. Lucente, *Dark Matter in the minimal Inverse Seesaw mechanism*, JCAP 10 (2014) 001; arXiv:1406.6556 [hep-ph].
- [17] D. J. Robinson and Y. Tsai *A Dynamical Framework for KeV Dirac Neutrino Warm Dark Matter*, Phys. Rev. **D 90**, 045030 (2014), arXiv:1404.7118 [hep-ph].
- [18] R. Krall, M. Reece and T. Roxlo, *Effective field theory and keV lines from dark matter*, arXiv:1403.1240 [hep-ph].
- [19] H. Ishida, K.S. Jeong and F. Takahashi, *7 keV sterile neutrino dark matter from split flavor mechanism*, arXiv:1402.5837 [hep-ph]
- [20] S. Patra and P. Pritimita, *7 keV sterile neutrino Dark Matter in extended seesaw framework*, arXiv:1409.3656 [hep-ph].
- [21] D. P. Finkbeiner and N. Weiner, *An X-Ray Line from eXciting Dark Matter*, arXiv:1402.6671 [hep-ph].
- [22] C. E. Aisati, T. Hambye, and T. Scarna, *Can a millicharged dark matter particle emit an observable gamma-ray line?*, arXiv:1403.1280 [hep-ph].
- [23] M. T. Frandsen, F. Sannino, I. M. Shoemaker, and O. Svendsen, *X-ray Lines from Dark Matter: The Good, The Bad, and The Unlikely*, JCAP 1405 (2014) 033, arXiv:1403.1570 [hep-ph].
- [24] R. Allahverdi, B. Dutta, and Y. Gao, *keV Photon Emission from Light Nonthermal Dark Matter*, Phys. Rev. **D 89**, 127305 (2014); arXiv:1403.5717 [hep-ph].
- [25] J. M. Cline, Y. Farzan, Z. Liu, G. D. Moore, and W. Xue, *3.5 keV X-rays as the 21 cm line of dark atoms, and a link to light sterile neutrinos*, Phys. Rev. **D 89**, 121302 (2014), arXiv:1404.3729 [hep-ph].
- [26] H. Okada and T. Toma, *The 3.55 keV X-ray Line Signal from Excited Dark Matter in Radiative Neutrino Model*, Phys. Lett. **B 737** (2014) 162-166; arXiv:1404.4795 [hep-ph].
- [27] H.M. Lee, *Magnetic dark matter for the X-ray line at 3.55 keV*, Phys. Lett. **B 738** (2014) 118-122; arXiv:1404.5446 [hep-ph].
- [28] T. Higaki, K. S. Jeong, and F. Takahashi, *The 7 keV axion dark matter and the X-ray line*

- signal*, Phys.Lett. **B 733** (2014) 2531, [arXiv:1402.6965].
- [29] J. Jaeckel, J. Redondo, and A. Ringwald, *A 3.55 keV hint for decaying axion-like particle dark matter*, Phys.Rev. **D89** (2014) 103511, [arXiv:1402.7335].
- [30] H. M. Lee, S. C. Park, and W.-I. Park, *Cluster X-ray line at 3.5 keV from axion-like dark matter*, Eur.Phys. J. **C 74** (2014) 9, 3062; arXiv:1403.0865 [astro-ph.CO].
- [31] M. Cicoli, J. P. Conlon, M. C. D. Marsh, and M. Rummel, *A 3.55 keV Photon Line and its Morphology from a 3.55 keV ALP Line*, Phys.Rev. **D 90**, 023540 (2014) ; arXiv:1403.2370 [hep-ph].
- [32] A. Dias, A. Machado, C. Nishi, A. Ringwald, and P. Vaudrevange, *The Quest for an Intermediate-Scale Accidental Axion and Further ALPs*, JHEP 1406 (2014) 037, arXiv:1403.5760 [hep-ph].
- [33] Joseph P. Conlon, Francesca V. Day, *3.55 keV photon lines from axion to photon conversion in the Milky Way and M31*, JCAP 11 (2014) 033; arXiv:1404.7741 [hep-ph].
- [34] M. Cicoli, J. P. Conlon, M. C. D. Marsh, and M. Rummel, *A 3.55 keV Photon Line and its Morphology from a 3.55 keV ALP Line*, Phys. Rev. **D 90**, 023540 (2014); arXiv:1403.2370 [hep-ph].
- [35] K. Kong, J.-C. Park and S. C. Park, *X-ray line signal from 7 keV axino dark matter decay*, Phys. Lett. **B 733** (2014) 217220, arXiv:1403.1536 [hep-ph].
- [36] K.-Y. Choi and O. Seto, *X-ray line signal from decaying axino warm dark matter*, Phys.Lett. **B 735** (2014) 92, [arXiv:1403.1782].
- [37] S. P. Liew, *Axino dark matter in light of an anomalous X-ray line*, JCAP 1405 (2014) 044, [arXiv:1403.6621].
- [38] C. Kolda and J. Unwin, *X-ray lines from R-parity violating decays of keV sparticles*, Phys. Rev. **D 90**, 023535 (2014), arXiv:1403.5580.
- [39] N. E. Bomark and L. Roszkowski, *The 3.5 keV X-ray line from decaying gravitino dark matter*, Phys. Rev. **D 90**, 011701 (2014) ; arXiv:1403.6503.
- [40] S. Demidov and D. Gorbunov, *SUSY in the sky or keV signature of sub-GeV gravitino dark matter*, arXiv:1404.1339.
- [41] K. Nakayama, F. Takahashi and T.T. Yanagida, *The 3.5 keV X-ray line signal from decaying moduli with low cutoff scale*, Phys. Lett. **B 735** (2014) 178-182 ,arXiv:1403.1733 [hep-ph].
- [42] K. Nakayama, F. Takahashi and T.T. Yanagida, *Anomaly-free flavor models for Nambu-*

- Goldstone bosons and the 3.5 keV X-ray line signal*, Phys. Lett. **B 734** (2014) 178-182, arXiv:1403.7390 [hep-ph].
- [43] F.S. Queiroz and K. Sinha, *The Poker Face of the Majoron Dark Matter Model: LUX to keV Line*, Phys. Lett. **B 735** (2014) 6974, arXiv:1404.1400 [hep-ph].
- [44] E. Dudas, L. Heurtier, and Y. Mambrini, *Generating X-ray lines from annihilating dark matter*, Phys. Rev. **D 90**, 035002 (2014); arXiv:1404.1927.
- [45] K. Babu and R. N. Mohapatra, *7 keV Scalar Dark Matter and the Anomalous Galactic X-ray Spectrum*, Phys. Rev. **D 89** (2014) 115011, arXiv:1404.2220 [hep-ph].
- [46] Y. Farzan and A.R. Akbarieh, *Decaying Vector Dark Matter as an Explanation for the 3.5 keV Line from Galaxy Clusters*, JCAP 11 (2014) 015; arXiv:1408.2950 [hep-ph].
- [47] A. Abada, V. De Romeri, A.M. Teixeira, *Effect of sterile states on lepton magnetic moments and neutrinoless double beta decay*, JHEP 1409 (2014) 074; arXiv:1406.6978 [hep-ph].
- [48] Z. Kang, P. Ko, T. Li and Y. Liu, *Natural X-ray Lines from the Low Scale Supersymmetry Breaking*, arXiv:1403.7742 [hep-ph].
- [49] A. Boyarsky, O. Ruchayskiy, and M. Shaposhnikov, *The role of sterile neutrinos in cosmology and astrophysics*, Ann.Rev.Nucl.Part.Sci.59:191-214,2009, arXiv:0901.0011 [hep-ph];
- [50] A. Kusenko, *Sterile neutrinos: the dark side of the light fermions*, Phys. Rept. 481:1-28,2009, arXiv:0906.2968 [hep-ph].
- [51] A. Adulpravitchai and M.A. Schmidt, *A Fresh Look at keV Sterile Neutrino Dark Matter from Frozen-In Scalars*, arXiv:1409.4330 [hep-ph].
- [52] J.M. Cline and A.R. Frey, *Consistency of dark matter interpretations of the 3.5 keV X-ray line*, arXiv:1410.7766 [astro-ph.CO]
- [53] Particle Data Group, J. Beringer *et al.*, Phys. Rev. **D 86**, 010001 (2012).
- [54] E. Ma, *Verifiable Radiative Seesaw Mechanism of Neutrino Mass and Dark Matter*, Phys. Rev. **D 73**, 077301 (2006), arXiv:0601225 [hep-ph].
- [55] E. Masso, S. Mohanty and S. Rao, *Dipolar Dark Matter*, Phys. Rev. **D 80**, 036009 (2009), arXiv:0906.1979 [hep-ph].
- [56] K. Griest and D. Seckel, *Three exceptions in the calculation of relic abundances*, Phys. Rev. **D 43**, 3191 (1991). A. Chatterjee and N. Sahu, *Resurrecting Sneutrino ($\tilde{\nu}_L$) Dark Matter in light of Neutrino Mass and LUX*, Phys. Rev. **D 90**, 095021 (2014); arXiv:1407.3030 [hep-ph].
- [57] D. S. Akerib *et al*, **LUX Collaboration**, *First results from the LUX dark matter experi-*

- ment at the Sanford Underground Research Facility*, Phys. Rev. Lett. **112**, 091303 (2014), arXiv:1310.8214 [astro-ph.CO].
- [58] J. Giedt, A. W. Thomas, and R. D. Young, *Dark matter, the CMSSM and lattice QCD*, Phys. Rev. Lett. **103**, 201802 (2009); arXiv:0907.4177 [hep-ph].
- [59] Y. Mambrini, *Higgs searches and singlet scalar dark matter: Combined constraints from XENON 100 and the LHC*, Phys. Rev. **D 84**, 115017 (2011), arXiv:1108.0671 [hep-ph].